

APPENDIX B: PLAN OF CARE — ASTHMA

STUDENT INFORMATION

School _____ Age _____ Homeroom Teacher _____ Student Name _____

PLAN OF CARE — ASTHMA

STUDENT INFORMATION

Student Name _____ Date Of Birth _____
 Ontario Ed. # _____ Age _____
 Grade _____ Teacher(s) _____

Student Photo (optional)

EMERGENCY CONTACTS (LIST IN PRIORITY)

NAME	RELATIONSHIP	DAYTIME PHONE	ALTERNATE PHONE
1.			
2.			
3.			

KNOWN ASTHMA TRIGGERS

CHECK (✓) ALL THOSE THAT APPLY

<input type="checkbox"/> Colds/Flu/Illness	<input type="checkbox"/> Change in Weather	<input type="checkbox"/> Pet Dander	<input type="checkbox"/> Strong Smells
<input type="checkbox"/> Smoke (e.g., tobacco, fire, cannabis, second-hand smoke)	<input type="checkbox"/> Mould	<input type="checkbox"/> Dust	<input type="checkbox"/> Cold Weather
<input type="checkbox"/> Physical Activity/Exercise	<input type="checkbox"/> Other (Specify) _____		
<input type="checkbox"/> At Risk for Anaphylaxis (Specify Allergen) _____			
<input type="checkbox"/> Asthma Trigger Avoidance Instructions: _____			
<input type="checkbox"/> Any Other Medical Condition or Allergy? _____			

APPENDIX B: PLAN OF CARE — ASTHMA

STUDENT INFORMATION

School _____

Age _____

Homeroom Teacher _____

Student Name _____

DAILY/ ROUTINE ASTHMA MANAGEMENT

RELIEVER INHALER USE AT SCHOOL AND DURING SCHOOL-RELATED ACTIVITIES

A reliever inhaler is a fast-acting medication (usually blue in colour) that is used when someone is having asthma symptoms. The reliever inhaler should be used:

When student is experiencing asthma symptoms (e.g., trouble breathing, coughing, wheezing).

Other (explain): _____

Use reliever inhaler _____ in the dose of _____
(Name of Medication) (Number of Puffs)

Spacer (valved holding chamber) provided? Yes No

Place a (✓) check mark beside the type of reliever inhaler that the student uses:

Airomir Ventolin Bricanyl Other (Specify) _____

Student requires assistance to **access** reliever inhaler. Inhaler must be **readily accessible**.

Reliever inhaler is kept:

With _____ – location: _____ Other Location: _____

In locker # _____ Locker Location: _____ Locker Combination: ____

Student **will carry** their reliever inhaler **at all times** including during recess, gym, outdoor, on the bus (if applicable) and during off-site activities.

Reliever inhaler is kept in the student's:

Pocket

Backpack/fanny Pack

Case/pouch

Other (specify): _____

Does student require assistance to **administer** reliever inhaler? Yes No

Student's **spare** reliever inhaler is kept:

In main office (specify location): _____ Other Location: _____

In locker #: _____ Locker Location: _____ Locker Combination: ____

CONTROLLER MEDICATION USE AT SCHOOL AND DURING SCHOOL-RELATED ACTIVITIES

Controller medications are taken regularly every day to control asthma. Usually, they are taken in the morning and at night, so generally not taken at school (unless the student will be participating in an overnight activity).

Use/administer _____ In the dose of _____ At the following times: _____
(Name of Medication)

Use/administer _____ In the dose of _____ At the following times: _____
(Name of Medication)

Use/administer _____ In the dose of _____ At the following times: _____
(Name of Medication)

APPENDIX B: PLAN OF CARE — ASTHMA

STUDENT INFORMATION

School _____

Age _____

Homeroom Teacher _____

Student Name _____

EMERGENCY PROCEDURES

IF ANY OF THE FOLLOWING OCCUR:

- Continuous coughing
- Trouble breathing
- Chest tightness
- Wheezing (whistling sound in chest)

(* Student may also be restless, irritable and/or quiet.)

TAKE ACTION:

STEP 1: Immediately use fast-acting reliever inhaler (usually a blue inhaler). Use a spacer if provided.

STEP 2: Check symptoms. Only return to normal activity when all symptoms are gone.

If symptoms get worse or do not improve within 10 minutes, this is an **EMERGENCY!**
Follow steps below.

IF ANY OF THE FOLLOWING OCCUR:

- Breathing is difficult and fast
- Cannot speak in full sentences
- Lips or nail beds are blue or grey
- Skin or neck or chest sucked in with each breath

(*Student may also be anxious, restless, and/or quiet.)

THIS IS AN EMERGENCY:

STEP 1: IMMEDIATELY USE ANY FAST-ACTING RELIEVER (USUALLY A BLUE INHALER). USE A SPACER IF PROVIDED.

Call 9-1-1 for an ambulance. Follow 9-1-1 communication protocol with emergency responders.

STEP 2: If symptoms continue, use reliever inhaler every 5-15 minutes until medical attention arrives.

While waiting for medical help to arrive:

- ✓ Have student sit up with arms resting on a table (do not have student lie down unless it is an anaphylactic reaction).
- ✓ Do not have the student breathe into a bag.
- ✓ Stay calm, reassure the student and stay by their side.
- ✓ Notify parent(s)/guardian(s) or emergency contact.

APPENDIX B: PLAN OF CARE — ASTHMA

STUDENT INFORMATION

School _____ Age _____ Homeroom Teacher _____ Student Name _____

HEALTHCARE PROVIDER INFORMATION (OPTIONAL)

Healthcare provider may include: Physician, Nurse Practitioner, Registered Nurse, Pharmacist, Respiratory Therapist, Certified Respiratory Educator, or Certified Asthma Educator.

Healthcare Provider's Name: _____

Profession/Role: _____

Signature: _____ Date: _____

Special Instructions/Notes/Prescription Labels:

If medication is prescribed, please include dosage, frequency and method of administration, dates for which the authorization to administer applies, and possible side effects.

*This information may remain on file if there are no changes to the student's medical condition.

AUTHORIZATION/PLAN REVIEW

INDIVIDUALS WITH WHOM THIS PLAN OF CARE (POC) IS TO BE SHARED:

Note: Only individuals involved in the daily/routine management require the entire Plan of Care. All others will receive Emergency Procedures Section only.

Please select one of the following:

DSNB Teaching and Support Staff, Niagara Student Transportation Services and food service providers.

Only those listed below: _____

Parent(s)/Guardian(s): _____ Date: _____

Signature

Student: _____ Date: _____

Signature

Principal: _____ Date: _____

Signature